



118 North Hospital Drive • Abbeville, LA 70510 • 337-893-5466 • www.abbevillegeneral.com

Abbeville General Referral & Criteria Form

(Referral and Treatment)

Patient Selection and Treatment Initiation

This section provides essential information on the emergency use authorization product combination of Casirivimab and Imdevimab for the treatment of mild to moderate COVID-19 in adults with positive results of direct SARS-CoV-2 viral testing who are 17 years of age and older weighing at least 40kg, and who are at higher risk for progressing to severe COVID-19 and/or hospitalization.

Referral can be made for any patient who is considered higher risk by criteria listed below or by consideration of benefit-risk for an individual patient by the referring healthcare provider. The patient must also have a positive COVID-19 test result within the past 24-48 hours.

Higher risk is defined as patients who meet at least one of the following criteria. Please select those that apply:

- Are \geq 65 years of age
- Have a body mass index (BMI) \geq 25
- Have chronic kidney disease
- Have diabetes
- Have immunosuppressive disease
- Are currently receiving immunosuppressive treatment
- Pregnancy
- HTN or Cardiovascular disease (including congenital heart disease)
- Chronic lung disease (for example, COPD, asthma [moderate to severe]. Interstitial lung disease, cystic fibrosis and pulmonary hypertension)
- Sickle Cell disease
- Individual patient benefit-risk: _____

Patient Name: _____ DOB: _____ Patient Phone #: _____

Referring Provider: _____

Please have Patient & Physician sign to validate that they have been informed about the risk and benefits and that they have been provided with the fact sheet:

Patient Signature _____

Date _____

Physician Signature _____

Date _____

Send referral, positive test result, and patient face sheet to:

ABBEVILLE GENERAL EUA Infusion Center Fax 337-892-6997

The patient will be contacted within 24-48 hours to schedule an appointment.

FOR AGH OFFICE USE ONLY: This Referral & Criteria form is to be scanned into the patients' chart in the EMR

- Schedule patient for Outpatient IV Infusion Therapy and Initiate COVID MED Casirivimab-Imdevimab Protocol.

Physician Signature _____

Date/Time _____

FACT SHEET FOR PATIENTS, PARENTS AND CAREGIVERS
EMERGENCY USE AUTHORIZATION (EUA) OF CASIRIVIMAB AND IMDEVIMAB FOR
CORONAVIRUS DISEASE 2019
(COVID-19)

You are being given a medicine called **casirivimab** and **imdevimab** for the treatment of coronavirus disease 2019 (COVID-19). This Fact Sheet contains information to help you understand the potential risks and potential benefits of taking casirivimab and imdevimab, which you may receive.

Receiving casirivimab and imdevimab may benefit certain people with COVID-19.

Read this Fact Sheet for information about casirivimab and imdevimab. Talk to your healthcare provider if you have questions. It is your choice to receive casirivimab and imdevimab or stop at any time.

WHAT IS COVID-19?

COVID-19 is caused by a virus called a coronavirus. People can get COVID-19 through contact with another person who has the virus.

COVID-19 illnesses have ranged from very mild (including some with no reported symptoms) to severe, including illness resulting in death. While information so far suggests that most COVID-19 illness is mild, serious illness can occur and may cause some of your other medical conditions to become worse. People of all ages with severe, long-lasting (chronic) medical conditions like heart disease, lung disease, and diabetes, for example, seem to be at higher risk of being hospitalized for COVID-19.

WHAT ARE THE SYMPTOMS OF COVID-19?

The symptoms of COVID-19 include fever, cough, and shortness of breath, which may appear 2 to 14 days after exposure. Serious illness including breathing problems can occur and may cause your other medical conditions to become worse.

WHAT IS CASIRIVIMAB AND IMDEVIMAB?

Casirivimab and imdevimab are investigational medicines used to treat mild to moderate symptoms of COVID-19 in non-hospitalized adults and adolescents (12 years of age and older who weigh at least 88 pounds (40 kg)), and who are at high risk for developing severe COVID-19 symptoms or the need for hospitalization. Casirivimab and imdevimab are investigational because they are still being studied. There is limited information known about the safety and effectiveness of using casirivimab and imdevimab to treat people with COVID-19.

The FDA has authorized the emergency use of casirivimab and imdevimab for the treatment of COVID-19 under an Emergency Use Authorization (EUA). For more information on EUA, see the “**What is an Emergency Use Authorization (EUA)?**” section at the end of this Fact Sheet.

WHAT SHOULD I TELL MY HEALTH CARE PROVIDER BEFORE I RECEIVE CASIRIVIMAB AND IMDEVIMAB?

Tell your healthcare provider about all of your medical conditions, including if you:

- Have any allergies

- Are pregnant or plan to become pregnant
- Are breastfeeding or plan to breastfeed
- Have any serious illnesses
- Are taking any medications (prescription, over-the-counter, vitamins, and herbal products)

HOW WILL I RECEIVE CASIRIVIMAB AND IMDEVIMAB?

- Casirivimab and imdevimab are two investigational medicines given together as a single intravenous infusion (through a vein) for at least 1 hour.
- You will receive one dose of casirivimab and imdevimab by intravenous infusion.

WHAT ARE THE IMPORTANT POSSIBLE SIDE EFFECTS OF CASIRIVIMAB AND IMDEVIMAB?

Possible side effects of casirivimab and imdevimab are:

- Allergic reactions. Allergic reactions can happen during and after infusion with casirivimab and imdevimab. Tell your healthcare provider or nurse, or get medical help right away if you get any of the following signs and symptoms of allergic reactions: fever, chills, low blood pressure, changes in your heartbeat, shortness of breath, wheezing, swelling of your lips, face, or throat, rash including hives, itching, headache, nausea, vomiting, sweating, muscle aches, dizziness and shivering.

The side effects of getting any medicine by vein may include brief pain, bleeding, bruising of the skin, soreness, swelling, and possible infection at the infusion site.

These are not all the possible side effects of casirivimab and imdevimab. Not a lot of people have been given casirivimab and imdevimab. Serious and unexpected side effects may happen. Casirivimab and imdevimab are still being studied so it is possible that all of the risks are not known at this time.

It is possible that casirivimab and imdevimab could interfere with your body's own ability to fight off a future infection of SARS-CoV-2. Similarly, casirivimab and imdevimab may reduce your body's immune response to a vaccine for SARS-CoV-2. Specific studies have not been conducted to address these possible risks. Talk to your healthcare provider if you have any questions.

WHAT OTHER TREATMENT CHOICES ARE THERE?

Like casirivimab and imdevimab, FDA may allow for the emergency use of other medicines to treat people with COVID-19. Go to <https://www.covid19treatmentguidelines.nih.gov/> for information on other medicines used to treat people with COVID-19.

It is your choice to be treated or not to be treated with casirivimab and imdevimab. Should you decide not to receive casirivimab and imdevimab or stop it at any time, it will not change your standard medical care.

WHAT IF I AM PREGNANT OR BREASTFEEDING?

There is limited experience treating pregnant women or breastfeeding mothers with casirivimab and imdevimab. For a mother and unborn baby, the benefit of receiving casirivimab and imdevimab may be

greater than the risk from the treatment. If you are pregnant or breastfeeding, discuss your options and specific situation with your healthcare provider.

HOW DO I REPORT SIDE EFFECTS WITH CASIRIVIMAB AND IMDEVIMAB?

Tell your healthcare provider right away if you have any side effect that bothers you or does not go away.

Report side effects to **FDA MedWatch** at www.fda.gov/medwatch or call 1-800-FDA-1088 or call 1-844-734-6643.

HOW CAN I LEARN MORE?

- Ask your health care provider.
- Visit www.REGENCOV2.com
- Visit <https://www.covid19treatmentguidelines.nih.gov/>
- Contact your local or state public health department.

WHAT IS AN EMERGENCY USE AUTHORIZATION (EUA)?

The United States FDA has made casirivimab and imdevimab available under an emergency access mechanism called an EUA. The EUA is supported by a Secretary of Health and Human Service (HHS) declaration that circumstances exist to justify the emergency use of drugs and biological products during the COVID-19 pandemic.

Casirivimab and imdevimab have not undergone the same type of review as an FDA-approved or cleared product. The FDA may issue an EUA when certain criteria are met, which includes that there are no adequate, approved, available alternatives. In addition, the FDA decision is based on the totality of scientific evidence available showing that it is reasonable to believe that the product meets certain criteria for safety, performance, and labeling and may be effective in treatment of patients during the COVID-19 pandemic. All of these criteria must be met to allow for the product to be used in the treatment of patients during the COVID-19 pandemic.

The EUA for casirivimab and imdevimab is in effect for the duration of the COVID-19 declaration justifying emergency use of these products, unless terminated or revoked (after which the products may no longer be used).

REGENERON

Manufactured by:
Regeneron Pharmaceuticals, Inc.
777 Old Saw Mill River Road
Tarrytown, NY 10591-6707

©2020 Regeneron Pharmaceuticals, Inc. All rights reserved.
Authorized: 11/2020